

Multi-GeV Neutrinos from Internal Dissipation in GRB Fireballs

P. Mészáros^{1,2} & M.J. Rees¹

¹Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, U.K.

²Dpt. of Astronomy & Astrophysics, Pennsylvania State University, University Park, PA 16803

ABSTRACT

Sub-photospheric internal shocks and transverse differences of the bulk Lorentz factor in relativistic fireball models of GRB lead to neutron diffusion relative to protons, resulting in inelastic nuclear collisions. This results in significant fluxes of $\nu_\mu(\bar{\nu}_\mu)$ of ~ 3 GeV and $\nu_e(\bar{\nu}_e)$ of ~ 2 GeV, scaling with the flow Lorentz factor $\eta < \eta_\pi \sim 400$. This extends significantly the parameter space for which neutrinos from inelastic collision are expected, which in the absence of the above effects requires values in excess of η_π . A model with sideways diffusion of neutrons from a slower wind into a fast jet can lead to production of $\nu_\mu(\bar{\nu}_\mu)$ and $\nu_e(\bar{\nu}_e)$ in the 2-25 GeV or higher range, depending on the value of η . The emission from either of these mechanisms from GRB at redshifts $z \sim 1$ may be detectable in suitably densely spaced detectors.

Subject headings: Gamma-rays: Bursts - Neutrinos - Cosmology: Miscellaneous

1. Introduction

Neutrinos in three different energy ranges are expected from gamma-ray burst sources. The most straightforward and also the hardest to detect are quasi-thermal 10-30 MeV neutrinos associated with the stellar collapse or merger event that triggers the burst (Kumar 1999, Ruffert & Jahnka 1999). The low energies imply an extremely low detection cross section, which for sources at cosmological distances makes the required detector volume prohibitive. At much higher energies, $\epsilon_\nu \gtrsim 2 \times 10^{14}$ eV neutrinos are expected from the photopion process in $p\gamma$ interactions between protons accelerated in shocks and the GRB photons produced by the same shocks associated with the fireball. These have a much higher cross section, and significant detection rates are predicted (Waxman & Bahcall 1997, Vietri 1998, Rachen & Mészáros 1998). Such neutrino events would directly test the shock acceleration mechanism, since they require relativistic protons with energies $\sim 10^{16}$ eV. A third type of neutrino emission, which is expected in the 5-15 GeV energy range (Derishev et al, 1999, Bahcall & Mészáros 2000), occurs only if neutrons are present

in the fireball, and the fluences are detectable in km³ detectors. These neutrinos arise from longitudinal decoupling of the n and p flows, and this is expected to occur whether or not shocks occur, provided a substantial fraction of neutrons is present in the fireball and the total fireball bulk Lorentz factor $\eta \gtrsim \eta_\pi \simeq 400$.

In this paper we point out that if internal shocks occur below the radiation photosphere, this can lead to a regime where rapid diffusion of neutrons both parallel and transverse to the radial direction occurs. This leads to neutrinos with energy $\epsilon_{\nu\bar{\nu}} \sim 3(\eta/\eta_{\pi,sh})(2/1+z)$ GeV for a range of η both below and above a critical value $\eta_{\pi,sh} \sim \eta_\pi^{4/5} \sim 120$. This extends considerably the parameter space available for neutrino production from np interactions, in contrast to the smooth homogeneous fireball case where $\eta \gtrsim \eta_\pi$ is required. Whereas in typical shocks responsible for MeV gamma-rays the neutrino efficiencies from pp are low due to increased coupling and lower optical depths (Paczynski & Xu 1994), here the neutrino fluence is maximized with np collisions in shocks at or below the pionosphere, where the inelastic optical depth is $\gtrsim 1$. The flux of such neutrinos may be detectable with km³ Cherenkov detectors currently under planning, for a sufficiently dense phototube spacing. Similar or higher neutrino energies and rates are also possible from bursts involving a slow wind and a fast jet, where neutrons from the slow wind diffuse sideways into the highly relativistic GRB beam.

2. Sub-photospheric shocks and n diffusion

In a relativistic GRB outflow consisting of $\gamma, e^\pm, p$ and n which is initially accelerating with a common bulk Lorentz factor $\Gamma \simeq r/r_o$, the dynamics is the same for a spherical geometry or for an outflow constrained within a channel of angular width $\theta_j \gtrsim \Gamma^{-1}$. There are two dynamical regimes, depending on whether $\eta = L/\dot{M}c^2$ is greater or less than the critical value

$$\eta_\pi = \left(\frac{L\sigma_\pi}{4\pi m_p c^3 r_o (1+\xi)} \right)^{1/4} \simeq 4.6 \times 10^2 L_{52}^{1/4} r_{o7}^{-1/4} (1+\xi)^{-1/4} \quad (1)$$

where $\xi = n_n/n_p$ is the neutron/proton ratio, which for simplicity is taken to be constant over the range of interest, $\sigma_\pi \sim 3 \times 10^{-26}$ cm² is the inelastic cross section near threshold, and $r_o = 10^7 r_{o7}$ cm is the initial radius of the outflow. The parameter η_π of equation (1) can also be written as $\eta_\pi = \eta_*(\sigma_\pi/\sigma_T)^{1/4}$, where σ_T is the Thomson cross section and

$$\eta_* = \left(\frac{L\sigma_T}{4\pi r_o m_p c^3 (1+\xi)} \right)^{1/4} \simeq 10^3 L_{52}^{1/4} r_{o7}^{-1/4} (1+\xi)^{-1/4} \quad (2)$$

is the critical value of η for γ, p decoupling in a p, n outflow, which enters also in the definition of the photospheric radius (e.g. Mészáros & Rees 2000 for a pure proton fireball).

For $\eta \geq \eta_\pi$, the radial velocity of the n and p becomes decoupled while both components are in the accelerating regime where $\Gamma \propto r$, leading to relative drift velocities between n and p of

$v_{rel} \rightarrow c$, and 5-10 GeV neutrinos are expected from inelastic np collisions (Bahcall & Mészáros , 2000). On the other hand, for $\eta < \eta_\pi$ a decoupling of the bulk velocity of the n and p in the radial direction is prevented by the fact that the radial motion of both the n and p saturate to an asymptotic Lorentz factor $\Gamma_p \sim \Gamma_n \sim \eta$ at $r \gtrsim r_s \simeq r_o \eta$ before a relative drift $v_{rel} \rightarrow c$ has been reached, and thereafter the various components coast together. If η is constant in time and uniform across the channel of angular width θ_j , the comoving temperature $T' = T_o(r_o/r)$ or $T' = (T_o/\eta)(\eta/r)^{2/3}$ for r/r_o below or above the value η , and $T_o = 1.2L_{52}^{1/4} r_{07}^{-1/2}$ MeV is the initial temperature at r_o (henceforth denoting with primes quantities measured in the comoving frame). The thermal velocities of the baryons are therefore subrelativistic and the relative CM kinetic energy of the n and p is below the threshold energy ~ 140 MeV required for inelastic nuclear collisions leading to neutrinos.

However, for $\eta \gtrsim \eta_\pi$, it is possible and likely that temporal or angular inhomogeneities in η do exist in realistic GRB outflows. Both types of inhomogeneities imply the presence of density gradients, and thus the possibility for neutrons to diffuse into regions whose bulk velocity is significantly different from that where they originated. When the neutrons encounter protons or other neutrons with relative velocities $v_{rel} \gtrsim 0.6c$ the CM inelastic threshold energy $\epsilon_{rel} \gtrsim 140$ MeV is reached, leading to pion formation and neutrinos. In order for this to occur with significant probability, one requires the inelastic np comoving mean-free path $\ell'_{np} \sim (n'_p \sigma_\pi)^{-1}$ to be comparable to the scale of the inhomogeneity in the comoving frame, where $n'_p = L/[4\pi r^2 m_p c^3 \Gamma \eta (1 + \xi)]$ is the comoving proton density (henceforth primed quantities are in the comoving frame). In the case of internal shocks for which $\Delta\Gamma \sim \Gamma$, which ensures $v_{rel} \rightarrow c$, the characteristic length scale of the shocks is r/Γ . In the case of sideways diffusion, a thermal velocity $v_t \sim v_{rel} \rightarrow c$ also requires internal shocks to have re-heated the gas, so both effects are tied to each other, and the transverse diffusion is limited to the causal angle Γ^{-1} so the lengthscale is again r/Γ . In the saturated regime (which is a prerequisite for internal shocks to occur), when the n, p thermal relative velocities $v_{rel} \rightarrow c$ so $\sigma \rightarrow \sigma_\pi$, the inelastic optical depth parallel or perpendicular to the shock normal becomes smaller than unity above a radius r_π given by

$$(r_\pi/r_o) = \eta_\pi (\eta_\pi/\eta)^3 . \quad (3)$$

In the above expression the η_π dependence includes a factor $(1 + \xi)^{-1/4}$, which assumes protons as the only targets, for comparison with the radial decoupling case (the case where both n and p are targets is given by the same equation (3) by formally setting $\xi = 0$ in the η_π factor). The internal shocks, required to provide $v_{rel} \rightarrow c$, occur in a coasting regime, at radii

$$r_{sh}/r_o \sim (t_v/t_o)\eta^2 , \quad (4)$$

where t_v is the observer-frame variability timescale on which η varies significantly, and $t_v \gtrsim t_o \sim r_o/c \sim 10^{-3}$ s, where t_o is the minimum dynamical timescale in the system over which such variation can occur involving a substantial fraction of the total outflow energy. This defines a minimum shock radius $r_{sh,m}/r_o \sim \eta^2$, for $\eta < \eta_*$. For radii $r_{sh,m} \lesssim r \lesssim r_\pi$ shocks leading to $v_{rel} \rightarrow c$ are possible, while the additional condition $\tau'_\pi \geq 1$ ensures efficient pion production and

neutrino emission. This region is shown by horizontal hashed lines in Figure 1. The requirement that $r_{sh,m} \lesssim r_\pi$ translates into the condition

$$\eta \leq \eta_{\pi,sh} \sim (t_o/t_v)^{1/5} \eta_\pi^{4/5} \simeq 120 L_{52}^{1/5} (t_o/t_v)^{1/5} , \quad (5)$$

For radii above r_π , or for values of $\eta \gtrsim \eta_{\pi,sh}$, shocks and inelastic collisions can also occur in the saturated regime at $r > r_{sh,m}$, but the mean-free path ℓ'_π is longer than the characteristic system length r/Γ , and the optical depth for inelastic collisions $\tau'_\pi < 1$, being equal to the inverse ratio of these two lengths. This regime is shown with vertical hashed lines in Figure 1. For $\eta > \eta_* > \eta_\pi$ the neutrons have decoupled at a terminal value $\Gamma_{nf} \sim (3/4)\eta_\pi(\eta_\pi/\eta)^{1/3}$ (Bahcall & Mészáros 2000, Derishev et al 1999), while the protons (which are the component capable of being heated by shocks) reach a maximum bulk Lorentz factor $\Gamma_{pf} \sim \eta_* < \eta$ (Mészáros & Rees 2000). Shocks can still occur, but the radius at which they occur reaches a maximum value $r_{sh,M}/r_o \sim (t_v/t_o)\eta_*^2 \simeq$ constant.

Below the radius r_π the pionization optical depth $\tau'_\pi \gtrsim 1$. The neutrino emission from inelastic np collisions at threshold leading to pion and muon decay contains, on average, one ν_μ and one $\bar{\nu}_\mu$ per neutron, with mean CM energies of 30-50 MeV, and also either a ν_e or $\bar{\nu}_e$ with an average CM energy of about 30 MeV, resulting from pions and muons decaying at rest in the CM (Bahcall & Mészáros, 2000). Typical conditions assumed for internal shocks involve collisions between shells with saturated Lorentz factors differing by factors of a few, e.g. a root-mean square value $\eta = \sqrt{\eta_A \eta_B} = 10^2 \eta_2$ with $\eta_A = 200$, $\eta_B = 50$ as an example. If the luminosity is constant between variations, this corresponds to a mechanical dissipation efficiency $\varepsilon = (\eta_A + \eta_B - 2\sqrt{\eta_A \eta_B})/(\eta_A + \eta_B) = 0.2$. The typical collision Lorentz factor is $\Gamma_{rel} \simeq (1/2)(\eta_A/\eta_B + \eta_B/\eta_A) \simeq 2.12$, and the total CM energy is $W = (2m_p^2 + 2m_p E_{rel})^{1/2} \sim 2.35$ GeV, whereas the CM threshold energy is $2m_p + m_\pi \sim 2.01$ GeV. Thus, at depths $\tau_\pi > 1$ each n heated by an individual shock can, on average, collide $k_\pi \sim 2$ times (and produce a pion each time) before its CM energy drops below threshold. The observer-frame energy of the $\nu_\mu \bar{\nu}_\mu$ ($\nu_e \bar{\nu}_e$) which would be predominant in the detector is $\sim 50(30)\alpha\eta/(1+z)$ MeV, where $\alpha \sim 1$ near threshold, or

$$\epsilon_{\nu_\mu \bar{\nu}_\mu} \sim 3 \alpha \left(\frac{\eta}{\eta_{\pi,sh}} \right) \left(\frac{2}{1+z} \right) \text{ GeV} , \quad \epsilon_{\nu_e \bar{\nu}_e} \sim 2 \alpha \left(\frac{\eta}{\eta_{\pi,sh}} \right) \left(\frac{2}{1+z} \right) \text{ GeV} , \quad (6)$$

where $\eta_{\pi,sh}$ is given in equation (5). The average total $\nu + \bar{\nu}$ CM energy per neutron at threshold is $\epsilon'_{\nu\bar{\nu}} \sim 100\alpha k_\pi$ MeV, which in the observer frame is $\epsilon_{\nu\bar{\nu}} \sim 0.1\alpha k_\pi \eta(1+z)^{-1}$ GeV. The average neutrino detection cross section averaged over ν and $\bar{\nu}$ is (Gaisser, 1990) $\sigma_{\nu\bar{\nu}} \sim 0.5 \times 10^{-38} (\epsilon_{\nu\bar{\nu}}/\text{GeV}) \text{ cm}^2$, which (per neutron) is $\bar{\sigma}_{\nu\bar{\nu}} \sim 0.6 \times 10^{-37} \alpha (k_\pi/2) (\eta/\eta_{\pi,sh}) (2/(1+z)) \text{ cm}^2$. The total number of neutrons involved in the shock and diffusion process, with a shock dissipation efficiency $\varepsilon = 0.2\varepsilon_{0.2}$, is

$$N_{n,sh} = \left(\frac{\xi}{1+\xi} \right) \left(\frac{\varepsilon E}{\eta m_p c^2} \right) \sim 5.5 \times 10^{52} \varepsilon_{0.2} E_{53} \left(\frac{\eta}{\eta_{\pi,sh}} \right)^{-1} \left(\frac{2\xi}{1+\xi} \right) . \quad (7)$$

Assuming a km^3 detector with $N_t \sim 10^{39} N_{t39}$ target protons, and a burst rate out to a Hubble radius of $10^3 \mathcal{R}_b$ per year, the number of events per year in the detector will be $R_{\nu\bar{\nu}} \sim (N_t/4\pi D^2) \mathcal{R}_b N_n \bar{\sigma}_{\nu\bar{\nu}}$, where D is the proper distance out to redshift z . For an Einstein-de Sitter universe with Hubble constant $H = 65 h_{65} \text{ km/s/Mpc}$ this is

$$R_{\nu\bar{\nu},sh} \sim 3 E_{53} \varepsilon_{0.2} \alpha \left(\frac{k_\pi}{2} \right) \left(\frac{2\xi}{1+\xi} \right) N_{t39} \mathcal{R}_{b3} h_{65}^2 \left(\frac{2 - \sqrt{2}}{1 + z - \sqrt{1+z}} \right)^2 \text{ year}^{-1}, \quad (8)$$

in coincidence with observed GRBs, and independent of the value of η for $\eta \lesssim \eta_{\pi,sh}$, which is the limit below which $\tau_\pi \gtrsim 1$. For values $\eta_{\pi,sh} \lesssim \eta \lesssim \eta_\pi$, the rate (8) must be multiplied by $\tau'_\pi \sim (\eta_{\pi,sh}/\eta)^5$, so between these limits $R_{\nu\bar{\nu}}$ drops steeply. However, for $\eta \gtrsim \eta_\pi \sim 400$, even in a laminar flow without any shocks one expects radial decoupling of the np leading to inelastic collisions, in which case the rate becomes a factor $(k_\pi \varepsilon_{sh})^{-1} \sim 2.5$ larger, or (Bahcall & Mészáros 2000) $R_{\nu\bar{\nu}} \sim 7/\text{year}$ for typical $\epsilon_{\nu_\mu} \sim \epsilon_{\bar{\nu}_\mu} \sim 10 \text{ GeV}$.

3. External neutron injection into the beam

If the progenitor is a collapsing massive star, a relativistic fireball from the collapsed central engine, probably in the form of a jet, must punch its way out through the stellar core and the envelope in order to become a ‘manifest’ GRB which emits observable γ -rays. The collision of fast baryons from the jet with quasistatic or slow neutrons or baryons from the rest of the star is a possible source of high energy neutrinos. We show here that two of the most straightforward scenarios have a very low neutrino efficiency, and then go on to discuss a third scenario where the efficiency is significantly higher.

The simplest scenario is that of a beam-dump provided by the outer envelope of a massive progenitor star, before the jet has punched through it. The fast neutrons of the jet would penetrate ahead of the protons participating in the bow shock, and collide inelastically with the upstream baryons at relative Lorentz factors $\sim \eta_j \sim 10^2 - 10^3$. However, a jet with such a Lorentz factor takes only $t_{cross} \sim r/c\eta^2 \sim 3 \times 10^{-2} r_{13} \eta_2^{-2} \text{ s}$ in the observer frame to reach an envelope radius $10^{13} r_{13} \text{ cm}$, and for a burst of observed duration $t_b = 30(2/1+z) \text{ s}$ only a fraction $(t_{cross}/t_b) \sim 2 \times 10^{-3} r_{13} \eta_2^{-2} (t_b/30)^{-1}$ of the jet neutrons will collide with envelope targets before the latter is punched through. Assuming that the observer is in the jet beam, the isotropic equivalent output of jet neutrons is $N_n = 3 \times 10^{53} E_{53} \eta_2^{-1} (2\xi/1+\xi)$. The jet neutrons have a source frame energy $\epsilon_j \sim 10^2 \eta_2 \text{ GeV}$, well above the inelastic threshold for envelope baryons. Assuming that the pions decay at rest in the CM, the maximum observer frame energy of the ν_μ or $\bar{\nu}_\mu$ is $\epsilon_{\nu_\mu \bar{\nu}_\mu} \sim 50 \alpha \eta / (1+z) \text{ MeV} = 2.5 \alpha \eta_2 (2/1+z) \text{ GeV}$. The average total $\nu\bar{\nu}$ energy per neutron is $\epsilon \sim 0.1 \alpha \zeta_\pi \eta / (1+z) \text{ GeV}$ and the total detection cross section per neutron is $\bar{\sigma}_{\nu\bar{\nu}} \sim 2.5 \times 10^{-38} \alpha \zeta_\pi \eta_2 (2/1+z) \text{ cm}^2$, where ζ_π is the pion multiplicity. The expected rate of events is $R_{\nu\bar{\nu},dump} \sim (N_t/4\pi D^2) \mathcal{R}_b \bar{\sigma}_{\nu\bar{\nu}} N_n (t_{cross}/t_b) \sim 1.5 \times 10^{-2} E_{53} \alpha \zeta_\pi \eta_2^{-2} r_{13} (30\text{s}/t_b) (2/1+z) \text{ year}^{-1}$, with the same detector and cosmological scalings as in equation (8). This is therefore too

low for km^3 detectors.

A second natural scenario involves transverse neutron diffusion into the beam from the core of the evolved progenitor star, and illustrates some of the factors affecting neutrino efficiency, as well as serving as a preliminary to the third, more efficient scenario discussed in the subsequent paragraph. As the progenitor collapse proceeds the core will be heated by a shock moving outwards with $v = \beta_s c$ to temperatures characteristic of the virial value near the central engine region, and it will be permeated by a diffuse bath of thermal neutrinos and gamma rays capable of dissociating the metals in the core. The free neutrons then diffuse via elastic nuclear collisions, into the jet channel. Taking a core neutron density $n_n = \xi_c \bar{n}_c$ where $\bar{n}_c = 3 \times 10^{26} m_c r_{10}^{-3} \text{ cm}^{-3}$ is the baryon density for a core of m_c solar masses, the neutron flux diffusing into the channel is $\phi_n \sim (v_t \ell / t)^{1/2} n_n \text{ cm}^{-2} \text{ s}^{-1}$ after a time t , where $\ell \sim (\bar{n}_c \sigma_{el})^{-1}$, and $\sigma_{el} \sim \sigma_\pi (c/v_t)$ is the elastic cross section, $v_t \sim \beta_s c$ being the neutron thermal velocity in the core. Using an effective transverse area of the channel of $A \sim \pi r^2 \theta_j$, the time-integrated fluence of neutrons into the channel after time t is $N_{n,\perp} \sim 2t\pi r^2 \theta_j \beta_s (\xi/1 + \xi) (c\bar{n}_c / \sigma_\pi t)^{1/2}$, which is $N_{n,\perp} \sim 10^{51} \theta_{-1} r_{10}^{1/2} (m_c/2)^{1/2} (\beta_s/3) (2\xi/1 + \xi) (t_b/30\text{s})^{1/2} (2/1 + z)^{1/2}$, using a core mass $m_c = 2M_\odot$, an observed duration $t_b = 30(2/1 + z) \text{ s}$ and a jet angle $\theta = 10^{-1} \theta_{-1}$ radians. This is comparable to the number of baryons going out through the jet, which is $N_b \theta^2 \sim 6 \times 10^{51} E_{53} \eta_2^{-1} \theta_{-1}^2$, where N_b is the isotropic equivalent output of jet baryons, $N_b = 6 \times 10^{53} E_{53} \eta_2^{-1}$. The neutrons diffusing into the channel collide against the ultrarelativistic baryons in the jet, and as above, the maximum observed energy of the ν_μ or $\bar{\nu}_\mu$ produced is $\epsilon_{\nu_\mu \bar{\nu}_\mu} \sim 2.5 \alpha \eta_2 (2/1 + z) \text{ GeV}$ and the average detection cross section per neutron is $\bar{\sigma}_{\nu\bar{\nu}} \sim 2.5 \times 10^{-38} \alpha \zeta_\pi \eta_2 (2/1 + z) \text{ cm}^2$. However, the transverse inelastic optical depth through the jet encountered by the diffusing slow neutrons is $\tau_{\pi,\perp} \sim n_n \sigma_\pi r \theta \sim 5 \times 10^4 L_{52} \eta_2^{-1} r_{10}^{-1} \theta_{-1}$. Thus only a fraction $\tau_{\pi,\perp}^{-1}$ of the jet baryons can produce 2.5 GeV neutrinos, since the slow neutrons cannot penetrate far into the jet. The number of neutrino events in a km^3 detector is then $R_{\nu\bar{\nu},\text{core}} \sim (N_t/4\pi D^2) \mathcal{R}_b \bar{\sigma}_\nu 2N_b \tau_{\pi,\perp}^{-1} \sim 1.5 \times 10^{-4} \eta_2 \alpha \zeta_\pi r_{10} \theta_{-1}^{-1} (2/1 + z)^2 \text{ year}^{-1}$, with detector and cosmological scalings as in equation (8). This is far too low, and it indicates the critical role played by the transverse optical depth.

A maximally efficient scenario based on lateral neutron injection can be obtained when the jet transverse inelastic depth is of order unity, so that neutrons diffusing inwards can interact with the entire fast beam. The transverse inelastic depth of the jet baryons in the lab frame is

$$\tau_{\pi\perp} = n_j \sigma_\pi r \theta \sim \eta_\pi^4 \eta^{-1} \theta (r_o/r), \quad (9)$$

and unit optical depth is achieved at $r/r_o \sim 4 \times 10^6 \theta_{-1} \eta_3^{-1} L_{52} r_{07}^{-1}$ or about $4 \times 10^{13} \text{ cm}$ for a jet with $\eta = 10^3 \eta_3 \sim 1$. If the jet is surrounded by a slower outflow, say with $\Gamma_s \sim 10$ (and $\Gamma_s \lesssim \theta_j^{-1}$), the slow outflow can reach a distance $4 \times 10^{13} \text{ cm}$ in an observer frame burst duration, $r/c\Gamma_s^2 \sim t_b \sim 30 \text{ s}$, and internal shocks in the slow outflow can heat the baryons in it. Inelastic collisions will heat the neutrons in the slow flow carried along from lower radii to thermal velocities $v_t \lesssim c$, and lead also to $\nu\bar{\nu}$, but their lab-frame energy will be very low, $\lesssim 0.25 \text{ GeV}$. The lab frame density in the slow outflow will be 10^2 times larger than in the jet (assuming the same energy per solid angle), and neutrons can diffuse into the jet from a depth in the slow outflow which is

one-tenth of the width of the jet in the time taken for a neutron to cross the jet, the amount of slow mass in this slow width being ten times that in the jet. If the pions and muons are produced at rest in the CM, using the CM neutrino energies ϵ' of the previous section the maximum observer frame energy of the ν or $\bar{\nu}_\mu$ from inelastic collisions with jet baryons is $\epsilon \sim \epsilon' \eta (1+z)^{-1}$

$$\epsilon_{\nu\bar{\nu}_\mu} \sim 25 \alpha \eta_3 \left(\frac{2}{1+z} \right) \text{ GeV} \quad , \quad \epsilon_{\nu_e\bar{\nu}_e} \sim 15 \alpha \eta_3 \left(\frac{2}{1+z} \right) \text{ GeV} \quad , \quad (10)$$

where $\alpha \sim 1$ near threshold. The total observer frame neutrino energy per neutron is $\epsilon_\nu + \epsilon_{\bar{\nu}} \sim 0.1 \alpha \zeta_\pi \eta / (1+z)$ GeV, where ζ_π is the pion multiplicity, and the average detection cross section averaged over ν and $\bar{\nu}$ is $\bar{\sigma}_{\nu\bar{\nu}} \sim 2.5 \times 10^{-37} \alpha \zeta_\pi \eta_3 (2/(1+z)) \text{ cm}^2$ per neutron. The (isotropic equivalent) baryon output in the jet is $N_b = E/\eta m_p c^2 = 6 \times 10^{52} E_{53} \eta_3^{-1}$, and the number of expected neutrino detection events is

$$R_{\nu\bar{\nu},j} \sim (N_t/4\pi D^2) \mathcal{R}_b \bar{\sigma}_\nu 2N_b \sim 15 E_{53} \alpha \zeta_\pi N_{t39} \mathcal{R}_{b3} h_{65}^2 \left(\frac{2 - \sqrt{2}}{1+z - \sqrt{1+z}} \right)^2 \text{ year}^{-1} . \quad (11)$$

This assumes that the maximum emission occurs at $\tau_{\pi\perp} \sim 1$, while for $\tau_{\pi\perp} \sim E_{53} \eta_3^{-1} \theta_{-1} r_{13.6}^{-1} (t_b/30\text{s})^{-1} (1+z/2) > 1$ the rate would be reduced by $\tau_{\pi\perp}^{-1}$. Note that if the transverse neutron input $N_{n\perp} \sim 2\pi r^2 \theta \xi \beta_s (ct \bar{n}_s / \sigma_\pi)^{1/2} \sim 10^{53} \xi \beta_s L_{52}^{1/8} \eta_{s1}^{-1/2} t_1^{1/2} r_{07}^{3/8} r_{13.6} \theta_{-1}$ is less than the number of baryons in the jet $N_b \theta^2 = 6 \times 10^{50} E_{53} \theta_{-1}^2 \eta_3^{-1}$, then $N_{n\perp}$ must be used in equation (11), instead of N_b ; this would be the case only if the wind neutron content is very small, $\xi/\theta_{-1} < 0.006 L_{52}^{7/8} t_1^{1/2} \eta_3^{-1} \eta_{s1}^{1/2} \beta_s^{-1} r_{07}^{-3/8} r_{13.6}^{-1}$.

The above example used $\eta = 10^3$, but the optimum efficiency $\tau_{\pi\perp} \simeq 1$ can also be obtained for other values of η in such a slow+fast jet model. The rate (11), $R_{\nu\bar{\nu},j} \sim 15/\text{year}$, is independent of the value of η , since η^{-1} in the number of ejected baryons cancels out with η in the neutrino detection cross section. The neutrino energies do scale with η , and the same rate of 15/year would be obtained, e.g. for $\eta_j = 10^2$ with $\epsilon_{\nu\bar{\nu}_\mu} \sim 2.5$ GeV, $\epsilon_{\nu_e\bar{\nu}_e} \sim 1.5$ GeV, or for $\eta_j = 10^4$ with $\epsilon_{\nu\bar{\nu}_\mu} \sim 250$ GeV, $\epsilon_{\nu_e\bar{\nu}_e} \sim 150$ GeV. The latter is an extreme value of η_j , but such energy neutrinos would be more easily detectable with first generation km^3 detectors, which could therefore provide significant constraints on the jet Lorentz factor.

4. Discussion

We have discussed two new mechanisms for producing significant fluences of 2-25 GeV neutrinos in GRB fireballs. The first of these involves internal shocks occurring at or below the pionsphere (which itself is below the γ -ray photosphere), where the optical depth to inelastic nuclear collisions is large. This mechanism requires either temporal or angular inhomogeneities (or both) in the relativistic outflow, which otherwise can be considered as a simple flow (i.e. characterized by a single mean energy per solid angle and a single mean Lorentz factor, aside from order unity variations in it). This mechanism can occur for values

of $\eta \lesssim \eta_\pi^{4/5} (t_o/t_v)^{1/5} \lesssim 120 L_{52}^{1/5} r_{o7}^{-1/5} (t_o/t_v)^{1/5} (2/[1+\xi])^{1/5}$. Thus, it considerably relaxes the η parameter space requirements where neutrinos in the multi-GeV range can be expected.

The second mechanism involves transverse diffusion of neutrons from a slower outflow into a fast jet, at a height where the transverse inelastic optical depth of the latter is close to unity for maximal efficiency. This also requires the slow outflow to be heated by internal shocks at or below its own pionosphere, and requires a two-zone outflow, characterized by two different values of the mean Lorentz factor (or alternatively of the energy per solid angle). This mechanism operates optimally at radii where the transverse optical depth is near unity, and predicts nominal event rates of $\lesssim 15/\text{year}$ independent of η and neutrino energies $\epsilon_{\nu_\mu \bar{\nu}_\mu} \sim 25 \eta_3 (2/[1+z]) \text{ GeV}$, $\epsilon_{\nu_e \bar{\nu}_e} \sim 15 \eta_3 (2/[1+z]) \text{ GeV}$ which scale with η , providing a diagnostic for the value of the jet Lorentz factor.

A feature of these two mechanisms is that they can operate even if the outflow has initially a low neutron fraction, since inelastic collisions with optical depth $\gtrsim 1$ tend to convert p into n and viceversa with approximately equal probability. They may therefore occur either in neutron star mergers or in massive stellar progenitor collapses. For km^3 detectors such as ICECUBE (Halzen, 1999), ANTARES (Feinstein 1998), BAIKAL (Beloplatikov et al 1997) or NESTOR (Trascatti et al 1998) with sufficiently dense phototube spacing, the predicted event rates are $R_{\nu\bar{\nu}} \sim 3 - 15/\text{year}$, which would be detectable above the background in coincidence with GRB flashes. However, since these two mechanisms only require sub-photospheric internal shocks (while longitudinal decoupling does not require any shocks), 2-25 GeV or higher energy neutrinos could be observed from a much wider range of stellar collapse or compact merger events in the Universe at $z > 1$, not necessarily associated with γ -ray flashes.

This research has been supported by NASA NAG5-2857, the Guggenheim Foundation, the Sackler Foundation and the Royal Society. We are grateful to John Bahcall for valuable comments and discussions.

REFERENCES

- Bahcall, JN & Mészáros, P, Phys.Rev.Lett, in press (2000) (hep-ph/0004019)
- Beloplatikov, I.A. et al, 1997, Astroparticle Phys, 7, 263
- Derishev, EV, Kocharovsky, VV & Kocharovsky, VI.V 1999, ApJ 521, 640
- Feinstein, F, in *5th Int'l Workshop "Topics in Astroparticle & Underground Physics" (IAUP97)*, Gran Sasso, Italy; eds. A. Bottini et al, Nucl. Phys B 70, 445
- Gaisser, T.K., 1990, *Cosmic Rays and Particle Physics* (Cambridge: Cambridge Univ. Press);
- Vogel, P, Phys.Rev D, 29, 1918

- Halzen, F 1999, 17th Intl conf. Weak Interactions and Neutrinos, Cape Town, South Africa(astro-ph/9908138)
- Kumar, P, 1999, ApJ, 523, L113.
- Mészáros , P. & Rees, M.J., ApJ, 530, 292 (2000)
- Paczynski, B & Xu, G., 1994, ApJ, 427, 708
- Rachen, J & Mészáros , P, 1998, Phys Rev D, 58, 123005
- Ruffert, M & Janka, H-Th, 1999, Astron.Ap 344, 573
- Trascatti, L, 1998, in *5th Int'l Workshop "Topics in Astroparticle & Underground Physics"* (IAUP97), Gran Sasso, Italy; eds. A. Bottini et al, Nucl. Phys B 70, 442
- Vietri, M, 1998, Phys Rev Lett. 80, 3690;
- Waxman, E & Bahcall, JN 1997, Phys Rev Lett. 78, 2292;




Fig. 1.— Radial distance vs. bulk Lorentz factor η parameter space, showing the pionosphere (pionization optical depth $\tau_\pi = 1$), and the photosphere (scattering optical depth $\tau_\gamma = 1$). Horizontal hashed regions indicate $\tau_\pi > 1$, this being absent above η_π (where n and p are elastically coupled, until decoupling and pionization occur at $\tau_\pi \gtrsim 1$). Vertical hashed regions, only shown for part of the range, indicate $\tau_\pi < 1$. The bulk Lorentz factor saturation radius r_s is common for n and p below η_π , while above η_π protons saturate at $r_s(p)$ and neutrons saturate (decouple) at $\tau_\pi = 1$. Internal shocks are possible above the line $r_{sh}(t_o)$, marked by diagonal hashes.